

Yellowstone National Park forms the core of the Greater Yellowstone Ecosystem (GYE)—and at 28,000 square miles, is one of the largest intact temperate-zone ecosystems on Earth today.

Each of Yellowstone National Park's separate parts—the hydrothermal features, the wildlife, the lakes, the Grand Canyon of the Yellowstone River, and the petrified trees—could easily stand alone as a national park. That they are all at one place is testimony to Greater Yellowstone's diversity and natural wealth.

## Geoecosystem

Geological characteristics form the foundation of an ecosystem. In Yellowstone, the interplay between volcanic, hydrothermal, and glacial processes and the distribution of flora and fauna are intricate and unique.

The topography of the land from southern Idaho northeast to Yellowstone results from millions of years of hotspot influence (*see Chapter 3*). Some scientists believe the Yellowstone Plateau itself is a result of uplift due to hotspot volcanism. Today's landforms even influence the weather, channeling westerly storm systems onto the plateau where they drop large amounts of snow.

The distribution of rocks and sediments in the park also influences the distributions of flora and fauna. The volcanic rhyolites and tuffs of the Yellowstone Caldera are rich in quartz and potassium feldspar, which form nutrient-poor soils. Thus, areas of the park underlain by rhyolites and tuffs generally are characterized by extensive stands of lodgepole pine, which are drought tolerant and have shallow roots that take advantage of the nutrients in the soil. In contrast, andesitic volcanic rocks that underlie the Absaroka Mountains are rich in calcium, magnesium, and iron. These minerals weather into soils that can store more water and provide better nutrients than rhyolitic soils. These soils support more vegetative growth, which adds organic matter and enriches the soil. You can see the result when you drive over Dunraven Pass or through other areas of the park with Absaroka rocks. They have a more diverse flora, including mixed forests interspersed with meadows. Lake sediments such as those underlying Hayden Valley, which were deposited during glacial periods, form clay soils that allow meadow communities to out-compete trees for water. The patches of lodgepole pines in Hayden Valley grow in areas of rhyolite rock outcrops.

## GYE BASICS

- 12–18 million acres; 18,750–28,125 square miles (*see map, next page, for why it varies*)
- States: Wyoming, Montana, Idaho
- Encompasses state lands, two national parks, portions of six national forests, three national wildlife refuges, Bureau of Land Management holdings, private and tribal lands
- Managed by state governments, federal government, tribal governments, and private individuals
- One of the largest elk herds in North America

- Largest free-roaming, wild herd of bison in U.S.
- One of two grizzly populations in contiguous U.S.
- Home to the rare wolverine and lynx

## In Yellowstone National Park:

- 61 mammals
- 319 bird species; 148 species nest here
- 16 fish species: 11 native, 5 non-native
- 10 reptiles and amphibians
- 12,000+ insect species, including 128 species of butterflies
- 1,100+ species of vascular plants



These whitebark pine grow in the andesitic soils on Mount Washburn.


## Greater Yellowstone Ecosystem

Because of the influence rock types have on plant distribution, some scientists theorize that geology also influences wildlife distributions and movement. Whitebark pine is an important food source for grizzly bears during autumn. The bears migrate to whitebark pine areas such as the andesitic volcanic terrain of Mt. Washburn. Grazing animals such as elk

and bison are found in the park's grasslands, which grow best in sedimentary soil of valleys such as Hayden and Lamar. And the many hydrothermal areas of the park, where grasses and other food remain uncovered, provide a haven for animals during winter.

### How big is the Greater Yellowstone Ecosystem?

*Sizes, boundaries, and descriptions of any ecosystem can vary—and the GYE is no exception. The park most often uses the two figures listed on the previous page, and most often uses the map shown here.*



## Biological Diversity

Biological diversity is one of the benchmarks measuring the health of an ecosystem. Biodiversity can be measured two ways: the number of different species (also called richness) and the abundance of each species (also called evenness). The diversity of animals within the Greater Yellowstone Ecosystem is as great as that found anywhere in the lower 48 states.

Significantly, Greater Yellowstone's natural diversity is still essentially intact. With the exception of the black-footed ferret, the region appears to have retained or restored its full historic complement of vertebrate wildlife species—something truly unique in the wildlands of the contiguous 48 states.

The extent of wildlife diversity is due in part to the different habitats found in the region, ranging from high alpine areas to sagebrush country, hydrothermal areas, forests, meadows, and other habitat types. All of these are connected, including linkages provided by streams and rivers that course through the changing elevations.

Other unique lifeforms are protected here, too. Various species of microorganisms are the living representatives of the primitive lifeforms now recognized as the beginnings of life on this planet. Cyanobacteria found in Yellowstone's hot springs are similar to the cyanobacteria that were among the first organisms capable of photosynthesis (the process by which plants use sunlight to convert carbon dioxide to oxygen and other byproducts). Because Earth's original atmosphere was anoxic (without oxygen), cyanobacteria's photosynthesis began to create an atmosphere on Earth that would eventually support plants and animals.

## Cycles and Processes

Cycles and processes are the building blocks in the foundation of any ecosystem. Photosynthesis, predation, decomposition, climate, and precipitation facilitate the flow of energy and raw materials. Living things absorb, transform, and circulate energy and raw materials and release them again. Cycles and



processes are the essential connections within the ecosystem.

Lifeforms are active at all levels. Microbes beneath Yellowstone Lake thrive in hydrothermal vents where they obtain energy from sulphur instead of the sun. Plants draw energy from the sun and cycle minerals such as carbon, sulphur, and nitrogen through the system. Herbivores, ranging from ephydrid flies to elk, feed on the plants and, in turn, provide food for predators like coyotes and hawks. Decomposers—bacteria, fungi, other microorganisms—link all that dies with all that is alive.

The ecosystem is constantly changing and evolving. A forest fire is one example of such an integral, dynamic process. Fires rejuvenate forests on a grand scale. Some species of plants survive the intense burning to resprout. The serotinous cones of lodgepole pines pop open in heat generated by fires, spreading millions of seeds on the forest floor. After fire sweeps through an area, mammals, birds, and insects quickly take advantage of the newly created habitats. Fires recycle and release nutrients and create dead trees or snags that serve a number of ecological functions, such as the addition of organic matter to the soil when the trees decompose (*see Chapter 5*).

*The Lamar Valley's thick grasses grow in sedimentary soil laid down by glaciers. This and other Yellowstone grasslands provide habitat for bison, elk, deer, pronghorn, coyote, wolf, grizzly and black bear, golden and bald eagles, ravens, osprey, and many other species.*

*For many, the sight of a wolf chasing an elk through a meadow is the highest symbol of wildness. That's a good start, but it's the meadow that is telling us the most about what wildness really means.*

—Paul Schullery

*For more information about wolf recovery and elk on the northern range, see Chapter 9.*

## Increasing Complexity In the Community

Many scientists consider the restoration of the wolf to Yellowstone to be the restoration of ecological completeness in the Greater Yellowstone Ecosystem. This region now contains every large wild mammal, predator or prey, that inhabited it when Europeans first arrived in North America. But the wolf is only one factor—albeit restored—in the extremely complex and dynamic community of wild Yellowstone.

For the visitor, this community's complexity has been highlighted primarily through the large predators and their prey species. This ecological “suite” of species provides a rare display of the dramatic pre-European conditions of wildlife in North America.

### *Intricate Layers*

In the few short years since the wolves were restored, scientists have discovered layers of complexity reaching far beyond the photogenic and obvious large mammals. The striking change in willow height in some northern range willow communities is seen by some researchers as evidence that the wolves, by affecting the habits of the elk, may also affect vegetation.

More subtle and far more ecologically tangled changes are also occurring. For example, the carcasses of elk, bison, and other large mammals each become ecosystems of their own. Researchers have identified at least 57 species of beetle associated with these ungulate carcasses on the northern range. Only one of those 57 species eats ungulate meat. All the rest prey on other small scavengers, especially the maggots of fly larvae and the grubs of other beetles' larvae. Others consume carcass byproducts such as microscopic fungal spores. In this very busy neighborhood, thousands of appetites interact until the carcass melts away and everybody moves on.

Thus the large predators point us toward the true richness, messiness, and subtlety of wild Yellowstone. For a wolf pack, an elk is dinner waiting to happen; for beetles, flies, and many other small animals, the elk is a village waiting to happen.

### *The Flexible Balance of Nature*

In some public circles, there was an expectation that wolves would restore a “balance” to park ecosystems, a term usually meant to suggest animal populations would become stable and unchanging at levels pleasing to humans. Instead, a more dynamic variability is present, which no doubt characterized this region's wildlife populations for millennia. The “balance of nature” has long been a doubtful concept; nature does have balances, but they are fluid rather than static, flexible rather than rigid.

Consider the northern Yellowstone elk herd, which has declined for the past several years. The recovery of the wolf has occurred simultaneously with increases in grizzly bear and mountain lion populations, increased human hunting of elk north of the park, and an extended drought. Computer models developed prior to wolf recovery predicted a decline in elk, but did not incorporate these other factors, and the decline has exceeded those predictions. Populations of prey species that share their habitat with more, rather than fewer species of predators are now thought to fluctuate around lower equilibria. The elk populations of Yellowstone will no doubt continue to adjust to all the pressures and opportunities they face, just as all their wild neighbors, large and small, will.

While some people delight in the chance to experience the new completeness of the Yellowstone ecosystem, others are alarmed and angered by the changes. But with so few places remaining on Earth where we can not only preserve but study such ecological completeness, there seems little doubt about the extraordinary educational, scientific, and even spiritual values of such a wild community.



## Winter in the Ecosystem

Bison can reach food beneath three feet of snow, as long as the snow is not solidified by melting and refreezing. A bison's hump is made of elongated vertebrae to which strong neck muscles are attached, which enable the animal to sweep its massive head from side to side.



### Winter in Yellowstone

Deep snow, cold temperatures, and short days characterize winter in the Greater Yellowstone Ecosystem, conditions to which plants and animals are adapted. For example, conifers retain their needles through the winter, which extends their ability to photosynthesize. Aspens and cottonwoods contain chlorophyll in their bark, enabling them to photosynthesize before they produce leaves.

#### Animal Behavioral Adaptations

- Red squirrels and beavers cache food before winter.
- Some birds roost with their heads tucked into their back feathers to prevent heat loss.
- Deer mice huddle together to stay warm.
- Ungulates like deer, elk, and bison sometimes follow each other through deep snow to save energy.
- Small mammals find insulation, protection from predators, and easier travel by living beneath the snow.
- Grouse roost overnight by burrowing into snow for insulation.
- Bison, elk, geese, and other animals find food and warmth in hydrothermal areas.

#### Animal Morphological/Physical Adaptations

- Mammals molt their fur in fall. Incoming guard hairs are longer and protect the underfur. Additional underfur grows each fall and consists of short, thick, often wavy hairs designed to trap air. A sebaceous (oil) gland, adjacent to each hair canal, secretes oil to waterproof the fur. Mammals have muscular control of their fur, fluffing it up to trap air when they are cold and sleeking it down to remove air when they are warm.
- River otters' fur has long guard hairs with interlocking spikes that protect the underfur, which is extremely wavy and dense to trap insulating air. Oil secreted from sebaceous glands prevents water from contacting the otters' skin. After emerging from water, they replace air in their fur by rolling in the snow and shaking their wet fur.

- Snowshoe hares, long-tailed weasels, and short-tailed weasels turn white for winter. White color provides camouflage but may have evolved primarily to keep these animals insulated as hollow white hairs contain air instead of pigment.
- Snowshoe hares have large feet to spread their weight over the snow; martens and lynx grow additional fur between their toes to give them effectively larger feet.
- Moose have special joints that allow them to swing their legs over snow rather than push through snow as elk do.
- Chickadees' half-inch-thick layer of feathers keeps them up to 100 degrees warmer than the ambient temperature.

#### Biochemical/physiological

- Mammals and waterfowl exhibit counter-current heat exchange in their limbs that enables them to stand in cold water: Cold temperatures cause surface blood vessels to constrict, shunting blood into deeper veins that lie close to arteries. Cooled blood returning from extremities is warmed by arterial blood traveling towards the extremities, conserving heat.
- At night, chickadees undergo regulated hypothermia. Their body temperature drops from 108°F to 88°F, which lessens the sharp gradient between the temperature of their bodies and the external temperature. This leads to a 23 percent decrease in the amount of fat burned each night.
- Chorus frogs tolerate freezing by becoming severely diuretic in response to cold temperatures and the formation of ice within their bodies. At this point the liver quickly converts glycogen to glucose, which enters the blood stream and serves as an anti-freeze. Within eight hours, blood sugar rises 200-fold. When a frog's internal ice content reaches 60–65 percent, the frog's heart and breathing stop. Within one hour of thawing, the frog's heart resumes beating.

#### Types of Snow

**Temperature Gradient Snow** or "depth hoar," forms through snow metamorphosis during cold air temperatures when water moves from warmer snow near the ground to colder snow near the surface. Snow crystals grow in size, forming sugar snow where small mammals burrow.

**Equitemperature Snow** forms as new crystals of snow become rounded and snowpack settles.

**Rime Frost** forms when super-cooled water droplets contact an object and freeze in place.

**Hoar Frost** forms when water vapor sublimates onto a surface. Formation of surface hoar occurs when night temperatures are very low.

## Ecosystem Management Challenges

Despite the size of the ecosystem, Greater Yellowstone's biodiversity is in jeopardy. Many of its plant and animal species are considered to be rare, threatened, endangered, or of special concern. This includes more than 100 plants, hundreds of invertebrates, at least six fish species, several species of amphibians, at least 20 bird species, and 18 species of mammals. The numbers are estimates because, even in this vital region, comprehensive inventories have not been completed. Carnivorous mammals—including the grizzly bear, wolverine, and lynx—represent more than half of the mammals in danger.

Habitat modification—beyond the levels of natural disturbance—poses a serious threat to both biodiversity and to ecosystem processes. Such modifications fragment habitats and isolate populations of plants and animals from each other, cutting them off from processes necessary for survival.

Ecosystem management is gaining support among conservationists and resource managers who recognize that most protected parks and reserves represent fragments of much larger ecosystems. Ecosystem management addresses the whole ecosystem, including preserving individual components and the relationships and linkages between them. Maintaining healthy, functioning ecosystems more effectively preserves species than do emergency measures to bring back threatened species from the brink of extinction.

Ecosystem management includes human activities. Development proposals are evaluated using methods such as “cumulative effects analysis,” which considers combined effects of all development—not just one activity—on an entire area.

In the past, GYE has been managed as individual units drawn along political lines. The result has been fragmented, inconsistent, and sometimes contradictory management. Since the 1980s, however, the ecosystem management approach has been gaining support. For example, the supervisors, superintendents, and regional officials for the two national parks and six of seven national forests in the ecosystem meet periodically as the Greater Yellowstone Coordinating Committee to discuss common issues and seek solutions.

GYE is included in the Yellowstone to Yukon Conservation Initiative, or “Y2Y.” More than 170 organizations, institutions, and foundations based in Canada and the United States are working together to ensure the long-term survival of wildlife in the Northern Rockies from the Greater Yellowstone Ecosystem to the Yukon Highlands—a distance of 1,900 miles. Ecosystem management on this scale is needed for wide-ranging wildlife species such as grizzly bears and wolves; Y2Y seeks to build and maintain a life-sustaining system of core reserves and connecting wildlife corridors. Existing national, state, and provincial parks and wilderness areas will anchor the system, while the creation of new protected areas and cooperation of land-owners will provide the additional reserves and corridors.

## For More Information

Despain, D. G. 1987. The two climates of Yellowstone National Park. *Biological Science Proceedings of The Montana Academy of Science*. 47:11–20.

Forrest, Louise. 1988. *Field Guide to Tracking Animals in Snow*. Harrisburg, PA: Stackpole Books.

Halfpenny, James C. and Roy D. Ozanne. 1989 *Winter: An Ecological Handbook*. Boulder: Johnson Books

Marchand, Peter J. 1996 *Life in the Cold*. University Press of New England.

Meagher, M. and D. B. Houston. 1998. *Yellowstone and the Biology of Time*. Norman: Univ. of Oklahoma Press.

Wilmers, et al. 2003. Resource dispersion and consumer dominance: scavenging at wolf- and hunter-killed carcasses in Greater Yellowstone. *Ecology Letters*. 6:996–1003.

Wilmers, et al. 2003. Trophic facilitation by introduced top predators: grey wolf subsidies to scavengers in YNP. *J Animal Ecology*. 72:909–916

## Additional Information from Yellowstone National Park

Yellowstone National Park website, [www.nps.gov/yell](http://www.nps.gov/yell), includes an array of park information about resources, science, recreation, and issues.

*Yellowstone Science*, published quarterly, reports on research and includes articles on natural and cultural resources. Free; available from the Yellowstone Center for Resources, in the Yellowstone Research Library, or online at [www.nps.gov/yell](http://www.nps.gov/yell).

*Yellowstone Today*, published seasonally and distributed at entrance gates and visitor centers, includes features on park resources such as hydrothermal features.

Area trail guides detail geology of major areas of the park. Available for a modest donation at Canyon, Fountain Paint Pot, Mammoth, Norris, Old Faithful, and West Thumb areas.

Site Bulletins, published as needed, provide more detailed information on park topics such as bison management, lake trout, grizzly bears, and wolves. Free; available upon request from visitor centers.